

SUPPLEMENTARY TABLE S2. PREVALENCE OF RECOMMENDATIONS FOR COMPLEMENTARY HEALTH APPROACHES  
AMONG OFFICE-BASED PHYSICIANS, BY PHYSICIAN'S MEDICAL SPECIALTY: UNITED STATES, 2012

	<i>General and family practice % (95% KG CI)</i>	<i>Internal medicine % (95% KG CI)</i>	<i>Pediatrics % (95% KG CI)</i>	<i>Obstetrics and gynecology % (95% KG CI)</i>	<i>Psychiatry % (95% KG CI)</i>	<i>Other specialty % (95% KG CI)</i>	<i>p</i>
Any CHA	75.5 (71.40–79.33)	64.8 (58.08–71.07)	49.0 (42.48–55.59)	59.5 (51.28–67.36)	62.7 (54.07–70.75)	39.9 (37.27–42.67)	0.0000
Massage therapy	52.6 (47.52–57.60)	44.0 (36.89–51.38)	22.8 (17.16–29.24)	37.7 (29.67–46.22)	27.4 (19.90–35.91)	19.1 (17.05–21.34)	0.0000
Chiropractic or osteopathic manipulation	54.0 (48.97–59.07)	39.9 (32.94–47.14)	20.9 (15.83–26.76)	29.1 (21.52–37.67)	12.4 (7.23–19.32)	16.3 (14.23–18.56)	0.0000
Herbs and nonvitamin supplements	40.9 (35.90–46.07)	36.2 (29.27–43.66)	27.2 (21.30–33.83)	31.1 (23.24–39.81)	29.5 (22.26–37.64)	17.1 (15.10–19.33)	0.0000
Yoga	35.3 (30.45–40.47)	40.3 (33.19–47.80)	19.8 (14.55–26.02)	33.9 (25.86–42.73)	44.9 (36.44–53.59)	15.3 (13.24–17.49)	0.0000
Acupuncture	36.4 (31.59–41.50)	28.7 (22.20–35.99)	13.1 (8.43–19.09)	26.1 (18.49–34.85)	27.7 (20.00–36.57)	16.1 (13.98–18.40)	0.0000
Mind–body therapies	27.8 (23.17–32.91)	25.1 (18.59–32.48)	17.0 (11.90–23.08)	27.8 (20.08–36.68)	53.1 (44.48–61.60)	12.2 (10.36–14.27)	0.0000

Any CHA includes the following: massage therapy, chiropractic or osteopathic manipulation, herbs and other nonvitamin supplements, yoga, acupuncture, mind–body therapies (such as guided imagery, meditation, and progressive muscle relaxation, not including prayer), biofeedback or hypnosis, homeopathic treatment, and naturopathic treatment. *p* Values are based on Wald Chi-square tests.

CHA, complementary health approach; CI, confidence interval; KG, Korn–Graubard.

Source: National Ambulatory Medical Care Survey Physician Induction Interview, 2012 (*n* = 5622).