



Treatment payed completely by other, %	5.4 (109)	9.4 (105)	- B	- B	- B	- B	- B
(n)							
Patient believes that the pain will get better 0-10 (No chance - Very likely), mean	8.1 (2.1)	8.4 (2.0)	7.5 (2.3)	7.9 (2.1)	7.13 (2.3)	8.0 (1.9)	8.28 (1.9)
(SD)							
Lives alone, % (n)	12.7 (258)	12.5 (140)	14.6 (43)	12.2 (75)	11.9 (21)	8.9 (10)	13.4 (44)
MPI cluster ID / DYS / AC, % (n)	22.2/41.8/36.1 (1566)	19.3/45.2/35.5 (825)	29.2/36.4/34.3 (236)	23.6/38.6/37.8 (505)	20.7/32.0/47.3 (150)	25.3/48.4/26.3 (95)	24.6/38.8/36.5 (260)
Pain severity (MPI) 0-6, mean (SD)	3.4 (1.2)	3.5 (1.3)	3.5 (1.1)	3.3 (1.2)	3.2 (1.3)	3.7 (1.1)	3.3 (1.1)
Interference (MPI) 0-6, mean (SD)	2.9 (1.4)	2.9 (1.4)	2.9 (1.3)	2.8 (1.3)	2.6 (1.3)	3.1 (1.2)	2.8 (1.4)
Life control (MPI) 0-6, mean (SD)	3.5 (1.2)	3.5 (1.2)	3.4 (1.1)	3.5 (1.1)	3.7 (1.0)	3.3 (1.1)	3.5 (1.1)
Affective distress (MPI) 0-6, mean (SD)	2.6 (1.4)	2.5 (1.5)	2.9 (1.4)	2.7 (1.4)	2.6 (1.4)	2.9 (1.3)	2.7 (1.3)
Support (MPI) 0-6, mean (SD)	4.0 (1.7)	4.1 (1.7)	3.7 (1.7)	3.9 (1.7)	3.7 (1.7)	4.1 (1.6)	4.0 (1.7)
Punishing responses (MPI) 0-6, mean (SD)	0.9 (1.1)	0.9 (1.1)	1.1 (1.2)	1.0 (1.2)	0.85 (1.0)	1.1 (1.2)	1.0 (1.3)
Solicitous responses (MPI) 0-6, mean (SD)	2.8 (1.4)	3.0 (1.4)	2.6 (1.4)	2.6 (1.4)	2.5 (1.3)	2.9 (1.5)	2.6 (1.4)
Distracting responses (MPI) 0-6, mean (SD)	2.9 (1.4)	2.9 (1.4)	2.8 (1.4)	2.8 (1.5)	2.8 (1.6)	2.8 (1.4)	2.8 (1.4)
Pain intensity at 1st visit (first measure) 0-10, mean (SD)	5.6 (2.2)	5.7 (2.3)	5.6 (1.4)	5.38 (2.1)	5.24 (2.1)	5.9 (2.0)	5.3 (2.1)
EQ5D score baseline, mean (SD)	0.67 (0.3)	0.66 (0.2)	0.66 (0.2)	0.69 (0.2)	0.70 (0.2)	0.64 (0.2)	0.69 (0.2)
Excellent	8.0 (163)	10.4 (117)	3.4 (10)	5.8 (36)	5.1 (9)	8.9 (10)	5.2 (17)

	Very good	32.1 (653)	33.4 (375)	29.8 (88)	30.8 (190)	35.2 (62)	25.0 (28)	30.5 (100)
Health in general (study start), % (n)	Good	34.6 (704)	31.5 (353)	37.6 (111)	39.0 (240)	39.8 (70)	33.9 (38)	40.2 (132)
	Quite poor	13.3 (271)	10.3 (116)	19.7 (58)	15.7 (97)	13.6 (24)	25.0 (28)	13.7 (45)
	Poor	3.0 (62)	2.5 (28)	4.7 (14)	3.2 (20)	4.0 (7)	1.8 (2)	3.4 (11)
	Missing	8.9 (180)	11.9 (133)	4.7 (14)	5.4 (33)	2.3 (4)	5.4 (6)	7.0 (23)
<b>4<sup>th</sup> visit</b>								
Chiropractor believes that MC is appropriate for patient, % (n)	-	-	-	78.9 (486)	69.3 (122)	81.3 (91)	84.8 (278)	
Has taken analgesic medication for the pain, % (n)	-	-	-	18.0 (111)	21.6 (38)	18.8 (21)	15.9 (52)	
Pain intensity at 4 <sup>th</sup> visit (second measure) 0-10, n = 246, mean (SD)	-	-	-	3.1 (2.1)	4.3 (2.2)	2.8 (1.9)	2.4 (1.7)	